

BULLYING IS A PAIN IN THE BRAIN Ebook

Bullying is a pain in the brain: Understanding the Psychological and Neurological Impact

Bullying has long been recognized as a serious social issue affecting individuals across all age groups, from schoolchildren to adults in the workplace. While its external effects—such as emotional distress, social withdrawal, and low self-esteem—are well-documented, recent research reveals that bullying also leaves deep imprints on the brain. The phrase "bullying is a pain in the brain" encapsulates the profound neurological and psychological impact bullying can have, highlighting the importance of understanding its effects on mental health and brain functioning.

In this comprehensive article, we'll explore the multifaceted effects of bullying on the brain, delve into the science behind these impacts, and discuss strategies for prevention and healing. Whether you're a parent, educator, mental health professional, or someone who has experienced bullying, understanding these dynamics is crucial for fostering healthier environments and supporting recovery.

How Bullying Affects the Brain: An Overview

Bullying isn't merely a social or emotional issue; it has tangible effects on the brain's structure and function. The chronic stress associated with bullying triggers biological responses that can alter brain development, especially in children and adolescents whose brains are still maturing.

Key points to understand include:

- The role of stress response systems
- Impact on brain regions associated with emotion and cognition
- Long-term neurological consequences

The Science Behind Bullying and Brain Impact

The Stress Response System and Cortisol Release

When an individual faces bullying, their body perceives it as a threat, activating the body's stress response system. The hypothalamic-pituitary-adrenal (HPA) axis releases cortisol, the primary stress hormone.

Effects of chronic cortisol elevation include:

- Impaired hippocampal function, affecting memory and learning
- Altered prefrontal cortex activity, impacting decision-making and impulse control
- Increased amygdala activity, heightening fear and anxiety responses

Prolonged exposure to stress hormones can cause structural changes in these brain regions, leading to lasting effects on mental health.

Alterations in Brain Structure and Function

Research utilizing neuroimaging techniques has shown that victims of bullying often exhibit:

- Reduced gray matter volume in areas responsible for emotional regulation, such as the prefrontal cortex
- Hyperactivity of the amygdala, leading to heightened fear and emotional responses
- Changes in the hippocampus, influencing memory formation and stress regulation

These alterations can predispose individuals to mental health disorders including depression, anxiety, and post-traumatic stress disorder (PTSD).

Psychological Effects Linked to Brain Changes

The neurological impacts of bullying translate into various psychological issues, such as:

- Anxiety and Panic Disorders: Heightened amygdala activity increases sensitivity to perceived threats.
- Depression: Changes in brain chemistry and structure can lead to feelings of hopelessness and low self-worth.
- Low Self-Esteem and Social Withdrawal: Damage to areas involved in self-perception and social cognition can make individuals feel unworthy or afraid to engage with others.
- Post-Traumatic Stress Disorder (PTSD): Repeated trauma from bullying can cause persistent flashbacks, hyperarousal, and emotional numbing.

Understanding these links emphasizes the importance of early intervention and psychological support for bullying victims.

Long-Term Neurological Consequences of Bullying

While some brain effects may be reversible with appropriate treatment and support, others can have lasting impacts.

Potential long-term consequences include:

- Persistent anxiety or mood disorders
- Difficulty in forming trusting relationships
- Impaired cognitive development, especially if bullying occurs during critical developmental periods
- Increased vulnerability to substance abuse or self-harm as coping mechanisms

Research indicates that early and effective intervention can mitigate many of these adverse outcomes.

Factors Influencing the Degree of Brain Impact

Not everyone reacts to bullying in the same way, and several factors determine the severity of neurological and psychological effects:

- Age: Children and adolescents are more susceptible due to ongoing brain development.
- Duration and Severity of Bullying: Chronic bullying causes more profound brain changes.
- Support Systems: Strong family, peer, and community support can buffer against adverse effects.
- Pre-existing Mental Health Conditions: Individuals with prior vulnerabilities may experience more intense impacts.
- Resilience and Coping Skills: Personal resilience can influence recovery trajectories.

Strategies to Mitigate the Brain Effects of Bullying

Addressing the neuropsychological impact of bullying requires a multifaceted approach:

Prevention and Education

- Implement anti-bullying programs in schools
- Promote awareness about the consequences of bullying
- Foster inclusive and supportive school cultures

Psychological Support and Therapy

- Cognitive-behavioral therapy (CBT) to reframe negative thought patterns

- Trauma-focused therapies for those with PTSD symptoms
- Mindfulness and stress reduction techniques to lower cortisol levels

Building Resilience and Coping Skills

- Encourage social skills development
- Promote self-esteem and confidence
- Teach emotional regulation strategies

Family and Community Involvement

- Provide family counseling and support
- Engage community organizations in bullying prevention efforts
- Establish safe spaces and helplines for victims

Healing the Brain: Hope and Recovery

While the effects of bullying on the brain can be serious, recovery is possible. Neuroplasticity, the brain's ability to reorganize itself, offers hope that with appropriate interventions, victims can recover and even strengthen neural pathways.

Key steps for healing include:

- Early identification and intervention
- Consistent psychological therapy
- Creating a safe and nurturing environment
- Encouraging positive social interactions
- Promoting physical health through exercise and nutrition

Research continues to explore innovative therapies, such as neurofeedback and mindfulness-based stress reduction, which have shown promise in restoring brain function affected by trauma.

Conclusion: Breaking the Cycle and Supporting Victims

Understanding that bullying is a pain in the brain underscores the importance of addressing both its psychological and neurological impacts. Combating bullying requires collective effort?from schools, families, healthcare providers, and communities?to create environments where individuals feel safe, supported, and empowered to heal.

By recognizing the profound effects bullying can have on brain development and mental health, we can better tailor prevention strategies and therapeutic interventions. Early action not only prevents long-term neurological damage but also fosters resilience and well-being for those affected.

Remember, recovery is possible, and with awareness, compassion, and appropriate support, victims of bullying can reclaim their mental and emotional health?moving from pain in the brain to peace and resilience.

Keywords: bullying, pain in the brain, psychological impact, neurological effects, stress response, brain structure, mental health, neuroplasticity, trauma, resilience, prevention strategies, therapy

Bullying is a pain in the brain ? a phrase that succinctly captures the profound impact that persistent bullying can have on mental health and cognitive function. While many are familiar with the emotional toll of bullying, recent research underscores that the effects extend deep into the brain, influencing how individuals think, feel, and relate to themselves and others. In this comprehensive guide, we explore the multifaceted ways in which bullying damages the brain, the underlying neuroscience, and strategies for healing and prevention.

Understanding the Connection: Bullying and Brain Health

Bullying isn't just a social issue; it's a neurological concern as well. When someone experiences repeated harassment or intimidation, their brain responds with a cascade of biological and psychological changes. These alterations can have immediate effects?such as heightened anxiety?and long-term consequences, including chronic stress, depression, and cognitive impairments.

The Psychological Impact of Bullying

Before diving into the neuroscience, it's essential to recognize the emotional toll:

- Anxiety and fear: Victims often develop chronic anxiety.
- Depression: Persistent feelings of worthlessness or sadness.
- Low self-esteem: Internalizing negative messages.
- Social withdrawal: Avoidance of interactions to escape harassment.

These psychological effects are rooted in biological changes within the brain, which can perpetuate a cycle of distress.

The Neuroscience of Bullying: How It Affects the Brain

The Stress Response and the Amygdala

The amygdala, often called the brain's "fear center," plays a central role in processing threats and emotional responses.

- Hyperactivation: Chronic bullying leads to an overactive amygdala, heightening sensitivity to perceived threats.
- Consequences: Increased anxiety, hypervigilance, and difficulty distinguishing real threats from perceived ones.

The Prefrontal Cortex and Executive Function

The prefrontal cortex (PFC) is responsible for decision-making, impulse control, and social behavior.

- Impairment: Ongoing stress from bullying can impair PFC functioning.
- Effects: Reduced ability to regulate emotions, poor judgment, and difficulty concentrating.

Hippocampus and Memory Formation

The hippocampus is critical for memory and learning.

- Impact of Stress: Elevated cortisol levels from prolonged stress can shrink hippocampal volume.
- Result: Memory problems, difficulty learning, and challenges in forming new associations.

The Long-Term Effects of Bullying on Brain Development

For children and adolescents, whose brains are still developing, the impact can be particularly severe:

- Altered neural pathways: Repeated exposure to bullying can reinforce negative thought patterns.
- Developmental delays: Impaired emotional regulation and social skills.
- Increased risk of mental health disorders: Such as anxiety disorders, depression, and even post-traumatic stress disorder (PTSD).

In adults, the effects might manifest as persistent anxiety, depression, or difficulties in social relationships, often rooted in earlier adverse experiences.

Biological Markers and Evidence

Research studies have provided concrete evidence linking bullying to brain changes:

- MRI scans showing reduced gray matter volume in key areas like the prefrontal cortex and hippocampus.
- Elevated cortisol levels indicating sustained stress.
- Altered connectivity between emotional regulation centers and cognitive control regions.

These findings highlight that bullying's impact isn't purely psychological but manifests physically within the brain's structure and function.

Strategies for Healing and Mitigating Brain Damage

While the effects of bullying can be profound, recovery and resilience are possible with appropriate interventions:

Psychological Therapies

- Cognitive Behavioral Therapy (CBT): Helps reframe negative thought patterns.
- Mindfulness and Stress Reduction: Reduce hyperactivation of the amygdala.
- Trauma-Informed Care: Address underlying trauma and promote healing.

Pharmacological Interventions

- Antidepressants: May help regulate neurotransmitter imbalances.
- Anti-anxiety medications: To manage hyperarousal symptoms.

Lifestyle and Supportive Measures

- Physical activity: Promotes neurogenesis and brain plasticity.
- Social support: Builds resilience and restores self-esteem.
- Healthy sleep patterns: Essential for brain recovery and emotional regulation.

Prevention and Building a Bully-Free Environment

Addressing the issue at its roots requires systemic change:

- School programs: Anti-bullying policies and awareness campaigns.
- Parental involvement: Teaching empathy and conflict resolution.
- Community efforts: Creating safe spaces and support networks.

Prevention not only reduces immediate harm but also minimizes long-term brain consequences for future generations.

Conclusion: Recognizing and Addressing the Brain Pain of Bullying

Bullying is a pain in the brain ? a vivid reminder that its impact extends beyond surface-level hurt feelings to deep structural and functional changes within the brain. Recognizing this connection underscores the importance of early intervention, comprehensive support, and prevention strategies. By understanding the neuroscience behind bullying, educators, parents, mental health professionals, and communities can work collaboratively to foster safe environments and promote healing for those affected. The journey from pain to resilience is challenging but achievable, with awareness, compassion, and targeted action guiding the way.

Remember: No one should suffer in silence. If you or someone you know is impacted by bullying, seek help from mental health professionals, trusted adults, or support groups. Healing the brain is possible, and every step toward a bully-free environment makes a difference.

QuestionAnswer What does the phrase 'bullying is a pain in the brain' mean? It emphasizes that bullying can cause mental and emotional distress, affecting a person's thoughts, feelings, and overall mental health. How does bullying impact mental health in children and teenagers? Bullying can lead to anxiety, depression, low self-esteem, and even suicidal thoughts, as it negatively affects their emotional well-being and brain development. What are some signs that someone is being mentally affected by bullying? Signs include withdrawal from friends, changes in mood or behavior, difficulty concentrating, declining academic performance, and expressions of hopelessness or sadness. What can schools and parents do to help children cope with the mental effects of bullying? They can promote open communication, provide counseling support, educate about bullying prevention, and foster a safe and inclusive environment to reduce its impact. Why is it important to address the mental 'pain in the brain' caused by bullying? Addressing this pain is crucial because ongoing emotional distress can lead to long-term mental health issues, affecting a person's overall well-being and quality of life.

Related keywords: bullying, mental health, emotional well-being, peer pressure, school violence, cyberbullying, self-esteem, psychological impact, adolescent health, bullying prevention

Pain A Very Short Introduction

pain a very short introduction

Pain is an intrinsic part of the human experience, serving both as a warning system and a complex phenomenon that influences physical, emotional, and psychological well-being. It can be acute or chronic, mild or severe, localized or widespread. Understanding pain's causes, mechanisms, and management is essential for improving quality of life. This comprehensive guide explores the multifaceted nature of pain, offering insights into its types, underlying mechanisms, diagnostic approaches, and treatment options.

Understanding Pain: An Overview

Pain is a vital biological function designed to alert the body to potential or actual injury. Without the sensation of pain, individuals might not recognize harmful stimuli, leading to further injury or health complications. However, pain is also a deeply personal experience, influenced by genetic, psychological, and environmental factors.

Types of Pain

Pain can be classified into several essential categories based on its duration, origin, and underlying mechanisms.

1. Acute Pain

- Definition: Short-term pain that arises suddenly due to injury, surgery, or illness.
- Characteristics: Sharp, localized, and usually resolves once the underlying cause is treated.
- Examples: Dental pain, cuts, burns, or postoperative discomfort.

2. Chronic Pain

- Definition: Persistent pain lasting longer than three to six months, often beyond the normal healing process.
- Characteristics: May be continuous or intermittent, often resistant to traditional treatments.
- Examples: Back pain, arthritis, neuropathy.

3. Nociceptive Pain

- Definition: Caused by damage or inflammation of tissues.
- Types:

- Somatic pain: Arises from skin, muscles, bones, or joints.
- Visceral pain: Originates from internal organs.
- Characteristics: Often described as aching or throbbing.

4. Neuropathic Pain

- Definition: Resulting from nerve damage or dysfunction.
- Characteristics: Burning, stabbing, or shooting sensations.
- Examples: Diabetes-related neuropathy, post-herpetic neuralgia.

5. Psychogenic Pain

- Definition: Pain influenced or exacerbated by psychological factors such as stress or emotional distress.
- Note: Often coexists with physical pain.

The Mechanisms Behind Pain

Understanding how pain works involves exploring the nervous system's complex pathways.

1. The Pain Pathway

- Nociceptors: Specialized nerve endings that detect harmful stimuli.
- Transmission: Nerve signals travel via the spinal cord to the brain.
- Perception: The brain interprets signals as pain.

2. Modulation of Pain

- The body has mechanisms to inhibit or amplify pain signals.
- Endogenous opioids (like endorphins) can suppress pain perception.
- Psychological factors, such as attention and emotion, influence pain experience.

3. Central Sensitization

- Chronic pain can lead to increased sensitivity in the nervous system.
- Resulting in heightened pain responses to normally non-painful stimuli.

Diagnosing Pain

Accurate diagnosis is crucial for effective management. Healthcare professionals rely on a combination of patient history, physical examinations, and diagnostic tests.

Key Components of Pain Assessment

- Location and radiation
- Intensity and duration
- Character and quality
- Aggravating and relieving factors
- Impact on daily life

Diagnostic Tools and Tests

- Imaging studies (X-ray, MRI, CT scans)
- Nerve conduction studies
- Laboratory tests
- Psychological assessments

Managing and Treating Pain

Effective pain management involves a multidisciplinary approach tailored to the individual's specific condition and needs.

Pharmacological Treatments

- **Analgesics:** Acetaminophen, NSAIDs
- **Opioids:** Morphine, oxycodone (used cautiously)
- **Adjuvant medications:** Antidepressants, anticonvulsants for neuropathic pain

Non-Pharmacological Therapies

- Physical therapy and exercise
- Psychological therapies (cognitive-behavioral therapy)
- Complementary medicine (acupuncture, massage)
- Biofeedback and relaxation techniques

Emerging and Advanced Treatments

- Neurostimulation (TENS, spinal cord stimulators)
- Interventional procedures (nerve blocks, injections)
- Regenerative medicine (stem cell therapy)

Chronic Pain and Its Impact

Chronic pain extends beyond physical discomfort, affecting emotional and social aspects of life.

- Psychological effects: Anxiety, depression, sleep disturbances.
- Social consequences: Reduced work productivity, strained relationships.
- Economic impact: Increased healthcare costs and lost income.

Preventing Pain and Promoting Wellness

Prevention strategies focus on lifestyle modifications and early intervention.

- Maintaining a healthy weight
- Engaging in regular physical activity
- Practicing good ergonomics
- Avoiding tobacco and excessive alcohol
- Managing stress through mindfulness and relaxation techniques

The Future of Pain Management

Research continues to advance understanding of pain, leading to innovative therapies:

- Personalized medicine approaches based on genetic profiles.
- Development of novel analgesics with fewer side effects.
- Use of virtual reality and digital health tools for distraction and pain relief.
- Enhanced understanding of the biopsychosocial model for holistic care.

Conclusion

Pain is a complex and multifaceted phenomenon that plays a crucial role in protecting the body but can also significantly diminish quality of life when chronic or poorly managed. Recognizing the

different types and mechanisms of pain allows for tailored treatment strategies, combining pharmacological, psychological, and physical therapies. As research progresses, new, more effective solutions continue to emerge, offering hope for improved pain management and enhanced well-being for millions worldwide.

By understanding the intricacies of pain, patients and healthcare providers can work together more effectively to overcome challenges, improve outcomes, and restore comfort and function.

Pain: An In-Depth Exploration of Its Complexity, Causes, and Management

Pain is an intrinsic aspect of the human experience, serving as both a vital warning system and a complex phenomenon that influences physical, psychological, and social well-being. Despite its universality, pain remains one of the most challenging symptoms to understand, diagnose, and treat effectively. This investigative review aims to delve into the multifaceted nature of pain, exploring its biological mechanisms, classifications, psychosocial factors, and current approaches to management. By synthesizing recent research and clinical insights, this article provides a comprehensive overview suitable for healthcare professionals, researchers, and informed readers seeking a deeper understanding of this pervasive phenomenon.

Understanding Pain: Definitions and Theoretical Frameworks

Pain is traditionally defined by the International Association for the Study of Pain (IASP) as "an unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage." This definition underscores the subjective and multifaceted nature of pain, encompassing not only physical sensations but also emotional responses.

Several theoretical models have been proposed to explain pain perception:

- Nociceptive Model: Focuses on the activation of nociceptors?specialized sensory neurons responding to potentially damaging stimuli.
- Neuropathic Model: Emphasizes abnormal processing of pain signals due to nerve injury or dysfunction.
- Central Sensitization Model: Describes increased responsiveness of neurons within the central nervous system, leading to heightened pain perception even in the absence of peripheral stimuli.
- Biopsychosocial Model: Recognizes the interplay of biological, psychological, and social factors influencing pain experience.

Understanding these frameworks is essential for appreciating the complexity of pain and tailoring effective treatment strategies.

The Biological Basis of Pain

Nociception and Signal Transmission

Pain begins at the periphery with nociceptors—free nerve endings located in skin, muscles, joints, and visceral organs. These specialized receptors respond to mechanical, thermal, or chemical stimuli indicative of potential tissue damage.

Once activated, nociceptors generate electrical signals transmitted via primary afferent fibers:

- A-delta fibers: Myelinated fibers responsible for quick, sharp, localized pain.
- C fibers: Unmyelinated fibers mediating dull, aching, and throbbing pain.

These signals travel to the dorsal horn of the spinal cord, where they undergo initial processing.

Central Nervous System Processing

From the spinal cord, pain signals ascend through the spinothalamic tract to various brain regions, including:

- Thalamus: Relay station for pain signals.
- Somatosensory cortex: Location and intensity perception.
- Limbic system: Emotional and affective components.
- Prefrontal cortex: Cognitive evaluation and modulation.

Neurotransmitters such as substance P, glutamate, serotonin, and norepinephrine modulate transmission at multiple levels, influencing pain perception and modulation.

Neuroplasticity and Chronic Pain

Persistent pain can induce neuroplastic changes—alterations in neural pathways and synaptic strength—that perpetuate pain even after tissue healing. Central sensitization, characterized by increased excitability of dorsal horn neurons, plays a significant role in chronic pain syndromes.

Classifications of Pain

Proper classification aids in diagnosis and treatment planning. The main categories include:

Acute vs. Chronic Pain

- Acute Pain: Short-term, typically protective, resolving with tissue healing.
- Chronic Pain: Persists beyond normal healing time, often defined as pain lasting longer than 3 to 6 months.

Physiological vs. Pathological Pain

- Physiological (Nociceptive) Pain: Normal response to harmful stimuli.
- Pathological Pain: Resulting from nerve injury or dysfunction, including neuropathic and centralized pain syndromes.

By Origin

- Somatic Pain: From skin, muscles, or joints; usually well-localized.
- Visceral Pain: From internal organs; often diffuse and poorly localized.
- Referred Pain: Perceived at a site different from the actual source.
- Radicular Pain: Due to nerve root compression or inflammation.

Specialized Categories

- Neuropathic Pain: Due to nerve damage or dysfunction (e.g., diabetic neuropathy).
- Nociplastic Pain: Resulting from altered pain modulation without evident tissue damage (e.g., fibromyalgia).

Psychosocial Factors Influencing Pain

Pain is not solely a biological phenomenon; psychological and social elements significantly influence its perception and impact.

Psychological Components

- Mood and Emotions: Anxiety, depression, and stress can amplify pain perception.
- Cognitive Factors: Catastrophizing, attention, and beliefs about pain affect coping strategies.
- Behavioral Responses: Avoidance, medication use, and activity levels influence outcomes.

Social and Cultural Influences

- Cultural norms shape pain expression and management.
- Social support can buffer or exacerbate pain experiences.
- Socioeconomic status affects access to care and resources.

Recognizing these factors is crucial for comprehensive pain management.

Current Approaches to Pain Management

Effective pain management requires a multimodal and individualized approach, integrating pharmacological, non-pharmacological, and psychosocial strategies.

Pharmacological Interventions

- Non-Opioid Analgesics: NSAIDs, acetaminophen.
- Adjuvant Medications: Antidepressants (e.g., amitriptyline), anticonvulsants (e.g., gabapentin).
- Opioids: Reserved for severe pain; risks include dependence and side effects.
- Topical Agents: Lidocaine patches, capsaicin.

Non-Pharmacological Techniques

- Physical therapy.
- Cognitive-behavioral therapy (CBT).
- Mindfulness and relaxation techniques.
- Interventional procedures (e.g., nerve blocks, neuromodulation).

Emerging and Future Therapies

- Targeted molecular therapies.
- Neuromodulation devices.
- Regenerative medicine approaches.
- Personalized pain medicine based on genetic and biomarker profiles.

Challenges and Future Directions in Pain Research

Despite advances, pain remains a significant clinical challenge. Key issues include:

- Subjectivity and Variability: Difficulty in quantifying pain.

- Chronic Pain Epidemic: Rising prevalence and socioeconomic burden.
- Opioid Crisis: Need for safer, effective alternatives.
- Understanding Central Mechanisms: Better insights into neuroplasticity and central sensitization.
- Personalized Medicine: Tailoring treatments based on individual genetics and psychosocial profiles.

Future research aims to develop objective biomarkers, improve pain assessment tools, and discover novel therapeutic targets.

Conclusion

Pain, as a complex and multifaceted phenomenon, encompasses biological, psychological, and social dimensions. Its diverse classifications and underlying mechanisms demand a comprehensive approach to diagnosis and management. Continued research into the neurobiological substrates, psychosocial influences, and innovative therapies holds promise for improving outcomes for millions suffering from acute and chronic pain. Recognizing pain's subjective nature and individual variability is essential for advancing personalized medicine and alleviating this pervasive human experience.

References

(In an actual publication, this section would include detailed citations of relevant scientific literature, guidelines, and review articles.)

Question What is pain in simple terms? Pain is an unpleasant sensation that signals potential or actual damage to the body, serving as a warning to protect ourselves. Why is pain important? Pain is important because it alerts us to injury or illness, helping us take action to prevent further harm. What are the different types of pain? Pain can be classified as acute (short-term) or chronic (long-term), and can also be categorized as nociceptive or neuropathic. How is pain measured? Pain is often measured using scales like the Visual Analog Scale (VAS) or Numeric Rating Scale (NRS), where patients rate their pain severity. What causes pain? Pain is caused by signals from nerve endings called nociceptors when tissues are damaged or threatened. Can pain be psychological? Yes, psychological factors like stress, anxiety, or depression can influence the perception and intensity of pain. How is pain managed? Pain management includes medications, physical therapy, psychological support, and sometimes surgery, tailored to the type and cause of pain. What is the difference between acute and chronic pain? Acute pain is sudden and short-lived, usually from injury, whereas chronic pain persists longer than three months and may have no clear cause. Why does pain sometimes persist without injury? Persistent pain without injury can result from nerve damage, inflammation, or conditions like fibromyalgia, where the nervous system amplifies pain signals. What advances are being made in pain research? Recent advances include new pain medications, better understanding of pain pathways, and therapies like neuromodulation

and personalized pain management approaches.

Related keywords: pain, introduction, brief, overview, sensation, nociception, discomfort, pathology, treatment, physiology

Read the full article online: [PAIN A VERY SHORT INTRODUCTION](#)